



The use of hand-raised psittacines for reintroduction: a case study of scarlet macaws (*Ara macao*) in Peru and Costa Rica

Donald Brightsmith^{a,*}, Jenifer Hilburn^{b,c,1}, Alvaro del Campo^{d,2}, Janice Boyd^e,
 Margot Frisius^c, Richard Frisius^c, Dennis Janik^b, Federico Guillen^b

^a Department of Biology, Duke University, Durham, NC 27712, USA

^b Fundación para la Restauración de la Naturaleza – Zoo Ave, La Garita, Alajuela, CR, USA

^c Amigos de las Aves Apartado 2306-4050, Alajuela, CR, USA

^d Rainforest Expeditions, Selva Reys, Aramburú 166 – 4B, Lima 18, Peru

^e Amigos de las Aves USA, 317 Thames Dr., Slidell, LA 70458, USA

Received 28 August 2003; received in revised form 8 May 2004; accepted 18 May 2004

Abstract

This study reports on three scarlet macaw (*Ara macao*) reintroduction projects using hand-raised birds in Peru and Costa Rica. The habitats at the release sites ranged from pristine tropical forest to forest fragments in an agricultural matrix. The combined first-year survival was 74% and the annual post first-year survival was 96%. Survival rates were very high despite a wide range in predator communities. Number of birds released explained 70% of the variation in survival with birds from larger releases having higher survival rates. Behavioral evidence suggests that birds established at the site facilitated survival of later releases. Breeding attempts were recorded at all three sites and hand-raised birds with wild mates successfully fledged young in Peru. Supplemental feeding post-release played an important role in keeping the birds near the release site and facilitating social interactions. This work shows that properly socialized hand-raised macaws can survive and breed in the wild but that ex-pets are not good release candidates.

© 2004 Published by Elsevier Ltd.

Keywords: Reintroduction; Macaw; Psittaciformes; Supplemental feeding; Peru; Costa Rica; Captive breeding; Disease; Hand-raised; Socialization

1. Introduction

Captive breeding and reintroduction are important management tools for endangered species (Balmford et al., 1996; Noss, 2001). However these projects are expensive, have a high risk of failure and are usually not properly documented, making it impossible to learn from the successes and mistakes of others (Griffith et al., 1989; Beck et al., 1994; Biggins et al., 1999). The role of captive breeding and reintroduction has been hotly debated especially in the field of parrot conservation in part because parrots are susceptible to various lethal, contagious diseases that may lie dormant for years

(Clubb, 1992; Derrickson and Snyder, 1992; Wiley et al., 1992; Balmford et al., 1996; Snyder et al., 1996, 1997; Gippoliti and Carpaneto, 1997). The family Psittacidae contains the highest proportion of species at risk of extinction of any large avian family yet many are kept and bred in captivity (Clubb and Clubb, 1992a; Johnson, 1992; Bennett and Owens, 1997; Collar, 1997). This provides many opportunities for reintroduction by private and public institutions (Clubb and Clubb, 1992b; Snyder et al., 1994; USFWS, 2002; Collazo et al., 2003; Juniper, 2003). Captive-raised animals usually perform poorly in comparison to wild-caught individuals but studies must continue to evaluate the potential of captive-raised birds because translocation is not an option when wild populations are endangered or extinct (Griffith et al., 1989; but see Sanz and Grajal, 1998; Collazo et al., 2003).

The scarlet macaw (*Ara macao*) is widely distributed throughout tropical America (Forshaw, 1989). The bird

* Corresponding author.

E-mail address: Djb4@duke.edu (D. Brightsmith).

¹ Present address: WCS Wildlife Survival Center, St. Catherines Island, 182 Camellia Road, Midway, GA 31324, USA.

² Present address: 843 S. Miller Street, Chicago, IL 60607, USA.

was formerly quite common but habitat loss, hunting and capture for pets have caused drastic declines and extinction in many areas, most notably Central America (Wiedenfeld, 1994; Juniper and Parr, 1998; Renton, 2000). This study compares three scarlet macaw release projects to document the techniques used and determine what factors correlate with high survival rates.

2. Study areas

Curú National Wildlife Refuge is a working farm located on the Nicoya Peninsula, in western Costa Rica (09°47'N, 84°56'W, elevation: sea level). It covers 1492 ha: 70% is natural forest and 30% human-created habitats (Schutt and Vaughan, 1995). Rainfall is strongly seasonal and totals 2000 mm per year. The site is a mix of tropical dry and tropical moist forest (Holdridge, 1967). There are no large raptors able to kill adult macaws. Wild scarlet macaws disappeared in the late 1960s.

The San Josecito Valley Center for Release is in a 200 ha valley approximately 16 km north of Golfito, Costa Rica (08°37'N, 83°15'W, elevation: sea level). Rainfall is aseasonal and totals about 6000 mm per year. This site will be referred to as Golfito. The valley floor is second growth forest ringed on three sides by low mountains covered in primary tropical wet forest. Golfo Dulce borders the fourth side. Adjacent to the valley is Piedras Blancas National Park (15,000 ha). Scarlet macaws were extirpated in the late 1950s (Janik et al., 2003). There are no large eagles at the site and *Spizatus* hawk-eagles occur at very low densities.

Tambopata Research Center (13°07'S, 69°36'W, elevation: 250 m) is located in SE Peru on the border between the Tambopata National Reserve (275,000 ha) and the Bahuaja-Sonene National Park (537,000 ha) over 20 km from the nearest permanent human settlement (Foster et al., 1994). Rainfall totals 3200 mm and is weakly seasonal (Brightsmith, in press). Primary tropical moist forest, *Guadua* bamboo patches and riparian successional forest of differing ages surround the site (Griscom and Ashton, 2003). The area has populations of large macaws (*Ara ararauna*, *A. chloroptera* and *A. macao*) and large raptors (*Harpia harpyja*, *Morphnus guianensis*, *Spizatus tyrannus*, *Spizatus ornatus* and *Spizastur melanoleuca*, Foster et al., 1994).

3. Methods

3.1. Rearing

Release candidates were captive-raised from native stock in Alajuela Costa Rica at Zoo Ave (Golfito) and Amigos de las Aves (Curú) or rescued from nests of wild birds (Tambopata, Table 1). Hand-raised birds were hatched in incubators or raised by their parents up to 2 weeks before being removed for hand raising. Seven Golfito birds were raised to fledging by their parents. At Amigos de las Aves no attempt was made to isolate the birds from casual human contact. At Zoo Ave the chicks were isolated from most human contact and visited only during feeding. The birds at both facilities were weaned off of hand feeding around 100 days when they were

Table 1
Summary of methods used in three scarlet macaw releases in Latin America

	Curú, CR	Golfito, CR	Tambopata, Peru
General			
Source of birds	Captive breeding	Captive breeding	Wild nests
Age at release	1.7–3.7 years ($x = 2.7$)	?	90–100 days
Pre-release methods			
In flight cages pre-release	Yes	Yes	No
Predator conditioning pre-release	No	No	No
Disease screening pre-release	Yes	Yes	No
Disease detected	^a	No	<i>Salmonella</i>
Kept with conspecifics during rearing	Yes	Yes	Yes
Feeding			
Hand fed pre-weaning	Yes	Yes	Yes
Hand fed post-weaning	No	No	Yes
Fed wild local foods pre-release	Yes	Yes	No
Supplemental feeding post-release	Yes	Yes	Yes
Interactions with people			
Isolated from contact during rearing	No	Yes	No
Given affection pre-fledging	Yes	No	Yes
Given affection post-fledging	No	No	Yes
Approach people post-release	No	No	Yes

^a See text for discussion of *Chlamydia* [Chlamydia] *psittaci* testing here.

112 placed in small flight cages and learned to feed them-
 113 selves. The birds were in groups at all times throughout
 114 raising. Post-weaning contact with humans was mini-
 115 mal. Five confiscated ex-pets were given to Zoo Ave and
 116 included in the releases. These birds were probably 2–5
 117 years old and were removed from the wild as chicks
 118 (Janik et al., 2003). All birds were marked with indi-
 119 vidually numbered metal leg bands.

120 In Tambopata younger chicks were removed for
 121 hand-raising from natural and artificial nests at age 5–15
 122 days from 1991 to 1993 (Nycander et al., 1995). In 1994,
 123 the second and third eggs were taken and incubated in
 124 the lodge. In all years, pairs of chicks were raised in
 125 small boxes (35 cm on a side) and not isolated from
 126 casual human contact.

127 3.2. Health screening and disease

128 Veterinarians conducted general fitness exams (Curú
 129 and Golfito), blood tests (Curú and Golfito) and general
 130 fecal exams (Golfito). All tests came up negative in
 131 Golfito (Janik et al., 2003). In Curú, all birds tested
 132 disease-free before transport to the release site, but after
 133 arrival one bird tested positive for *Chlamydiophila*
 134 [*Chlamydia*] *psittaci*. The bird was sacrificed and a nec-
 135 roscopy showed no evidence of disease. The bird was one
 136 that had not been raised in the Amigos facility. The
 137 result may have been a false positive or indicate expo-
 138 sure to *Chlamydiophila* before being acquired. In Tam-
 139 bopata no pre-release health screenings were performed.
 140 In 1994 researchers found 7 of 17 (41%) hand-raised
 141 birds tested positive for *Salmonella* but none of the wild
 142 birds did. Karesh et al. (1997) conclude that the source
 143 of the infection was live and dead chickens used to feed
 144 the researchers and visitors.

145 3.3. Pre-release training

146 At both Curú and Golfito macaws were held in avi-
 147 aries at the release site for at least 6 months. The birds
 148 were fed a mixture of basic diet (fruits, rice, beans, dog
 149 food, etc.) and wild foods. At Tambopata the birds re-
 150 ceived little pre-release training. Formal predator aver-
 151 sion training was not done at any of the sites however
 152 two Golfito birds were killed in the pre-release cage by a
 153 *Leopardus pardalis*, which made the survivors wary of
 154 terrestrial mammals.

155 3.4. Releases

156 At Curú pairs and trios of birds were released over 17
 157 days starting on 7 January 1999. Most birds left the
 158 immediate area upon release but returned within 1–5
 159 days. At Golfito birds were released on 14 different dates
 160 from May 1999 to December 2001. At Tambopata birds
 161 were not held in cages and releases consisted of indi-

162 vidual fledging age birds (80–100 days) flying in to the
 163 forest. Tambopata birds took 12 h to 3 days to return to
 164 the lodge to be fed (AC).

3.5. Survey techniques

166 At all three sites, supplemental feeding post-fledging
 167 played a vital role in surveying the populations. Birds
 168 received a standard diet similar to what they were raised
 169 on for 2 months (Curú), and 10 months (Tambopata)
 170 post-release. At Golfito standard diet items were avail-
 171 able continuously due to the regularly spaced release
 172 events. By the end of the first year the birds obtained
 173 nearly all their calories from wild foods, but supple-
 174 mental feeding of a few highly preferred items (sun-
 175 flower seeds in Curú and Golfito, crackers and bananas
 176 in Tambopata) continued throughout the study.

177 At Curú researchers did not identify individual birds
 178 but counted the total number of individuals on a daily
 179 basis. In Golfito birds were marked for individual
 180 identification using black ink on the bill, small cuts in
 181 the tail feathers and radio collars (Janik et al., 2003).
 182 Birds were censused daily at feeding stations and op-
 183 portunistically at nest boxes. Eighteen of 38 birds (47%)
 184 were equipped with radio collars (Holohil, model AI-
 185 2C, Bjork and Powell, 1995). These collared birds were
 186 monitored daily for the first two weeks post-release,
 187 once a week for the first 3 months and irregularly
 188 thereafter. This study reports on sightings at Golfito
 189 through December 2002.

190 In Tambopata, birds were marked with leg bands that
 191 could not be read from a distance. Individuals were
 192 identified at the lodge and at nests during January–May
 193 1994 (AC), February–March 1998 (AC), May–August
 194 1998 (AC), September 1999 (DB), and November 1999–
 195 March 2000, 2001, 2002 (DB and assistants). Re-
 196 searchers monitored seven natural and 12–17 artificial
 197 macaw nest sites during November–March each year
 198 from 1999 to 2002.

199 Birds that dispersed away from the release site and
 200 did not return were classified as mortalities. While this
 201 method obviously underestimates survival it is appro-
 202 priate where the goal of reintroduction is to establish a
 203 new population.

3.6. Data analysis

205 We calculated mean and variance of daily survival
 206 rates for released macaws following Mayfield (1975) and
 207 tested differences in survival rates using a Z-test (Hensler
 208 and Nichols, 1981). The *P*-values of these tests were
 209 corrected using a sequential Bonferroni analysis with
 210 overall $\alpha = 0.05$ (Sokal and Rohlf, 1995). All birds re-
 211 leased in the same month were considered part of the
 212 same release. Our data violated the assumption that
 213 survival probabilities were independent among individ-

uals so we used a grouped logistic regression where the release was the basic unit of analysis (Cox and Snell, 1989). The variables number released, number established at the site before the release, and site were included as independent variables in the model and the Mayfield first year percent survival was used as the dependent variable. Annual Mayfield survival rates were calculated as (mean daily survival rate)³⁶⁴. Variables that did not contribute significantly to the model were eliminated and the analysis rerun. Since proportions become unstable and highly variable with small denominators, we excluded releases with <3 birds from the regression analysis (Pyle et al., 1993). Data are presented as mean (x) \pm standard deviation (SD).

4. Results

4.1. Survival

Seventy one scarlet macaws were released and the overall survival rate was 89% per year. At Curú 10 of the 13 birds (77%) were still alive 4 years after release. At Golfito 34 birds were released including 22 hand-raised, seven parent-raised and five confiscated ex-pets. Four birds were released, captured and released again resulting in 38 release events. Of the 38 releases, 63% were alive and returning to the release site through December 2002. At Tambopata a total of 20 scarlet macaws were released from 1992 to 1995 (release age 80–100 days). Of these 55% were still alive as of March 2002.

For all birds combined, the first-year survival rate was 74% and the annual post first-year survival rate was 96% (Table 2). The overall daily survival of birds at Golfito was lower than the daily survival rate for Curú ($x_{\text{Golfito}} = 0.999319 \pm 1.8 \times 10^{-4}$, $x_{\text{Curú}} = 0.999825 \pm 1.01 \times 10^{-4}$, $Z = 2.43$, $P_{(\text{Bonferroni corrected})} = 0.03$) and Tambopata ($x_{\text{Tambopata}} = 0.999815 \pm 6.5 \times 10^{-5}$, $Z = 2.57$, $P_{(\text{Bonferroni corrected})} = 0.03$). First year daily survival rates at Golfito were significantly lower than at Curú ($x_{\text{Golfito}} = 0.998624 \pm 3.8 \times 10^{-4}$, $x_{\text{Curú}} = 0.999773 \pm 2.27 \times 10^{-4}$, $Z = 2.59$, $P_{(\text{Bonferroni corrected})} = 0.03$). First-year daily survival rates did not differ significantly between Golfito and Tambopata ($x_{\text{Tambopata}} = 0.999529 \pm 2.7 \times 10^{-4}$, $Z = 1.93$, $P_{(\text{Bonferroni corrected})} = 0.104$). Post

first-year daily survival rates did not vary significantly among sites (Table 2).

A total of 11 independent releases were conducted ranging in size from 1 to 13 birds. Due to small sample sizes (<3 birds) three releases were eliminated from the statistical analyses. The Mayfield first-year survival of released macaws was positively correlated with the number of birds released (Fig. 1, grouped logistic regression: $n = 8$ releases, $r^2 = 69.1$, $\chi^2 = 9.15$, $v = 1$, $P = 0.0025$). The field site (Golfito, Curú or Tambopata) did not contribute significantly to the model (grouped multiple logistic regression: $\chi^2 = 3.2$, $v = 2$, $P = 0.2$), nor did the number of released birds previously established at the site (grouped multiple logistic regression: $\chi^2 = 0.4$, $v = 1$, $P > 0.5$).

4.2. Breeding behavior

At all three sites birds have formed pairs: Curú 3 pairs, Golfito 5 pairs and Tambopata all 11 surviving hand-raised birds have wild mates. Nest boxes at all sites have been investigated by released macaws and used at both Golfito and Tambopata. At Golfito five different pairs have defended artificial nests and at least one pair laid eggs. At Tambopata hand-raised birds and their wild mates nested successfully. In total six such pairs have defended nests, five laid eggs, and three fledged a total of four chicks. At Curú pairs have apparently attempted to nest in natural tree cavities in two different years but no chicks have been produced. Researchers have not monitored nests at Curú or Golfito so causes of nest failure there are unknown. At Curú and Golfito it was not known which individuals attempted breeding. The approximate ages for first breeding attempts at all sites are Curú 4–7 years, Golfito 5–6 years (Janik et al., 2003), and Tambopata 7.2 ± 0.8 years ($n = 4$).

4.3. Reactions to humans/habituation

Released birds showed little fear of humans. At Curú, birds could be approached within 4–5 m (DB pers. obs.) while at Golfito birds would allow people within 8 m (Janik et al., 2003). No birds at Golfito or Curú approached people in search of food (Table 1). At Tam-

Table 2
Macaw survival at three release sites^a

	Released	First-year survival (%)	Annual survival post first-year (%)	Mortalities post first-year	Years monitored
Curú	13	92	96	2	4.2
Golfito	31	60	98	1	2.7
Tambopata	20	84	96	5	10
Total	64	74	96	8	16.9

Mortalities are the number of birds that died or disappeared >1 year after release. Birds that left the release area and were never seen again were considered mortalities. The years monitored indicates the number of years the birds were monitored starting with the release of the first bird.

^aSurvival rates are calculated using the Mayfield method (Mayfield, 1975).




Fig. 1. First year survival of scarlet macaws (*Ara macao*) in relation to the number of birds released (grouped logistic regression: $n = 8$ releases, $r^2 = 69.1$, $\chi^2 = 9.15$, $v = 1$, $P = 0.0025$). Three releases of <3 birds were not included in the analysis. First year survival was calculated using the Mayfield (1975) method. The curves show the fitted values and 95% confidence interval from the regression analysis.

296 bopata birds had no fear of humans and regularly ap-
297 proached people for food.

298 Three of the five ex-pets released in Golfito associated
299 more closely with humans. Although no quantitative
300 data exist, they apparently socialized less with the other
301 released macaws, strayed less from the immediate re-
302 lease area, perched lower in the trees near the staff area
303 and occasionally walked on the ground. Despite these
304 apparently maladaptive behaviors, all 5 ex-pets survived
305 at least 2 years post-release.

306 5. Discussion

307 5.1. Survival

308 This study shows that hand-raised scarlet macaws can
309 survive in the wild in a range of abiotic and biotic
310 conditions and that larger releases were more successful
311 than smaller ones. Survival in Golfito was lower than at
312 the other two sites apparently due to the small average
313 release size and the delay in establishing a core flock.
314 The first-year survival of reintroduced parrots is usually
315 well below 50% so our 74% survival was greater than
316 expected (Snyder et al., 1987, 1994; USFWS, 2002;
317 Collazo et al., 2003; but see Sanz and Grajal, 1998). The
318 high first-year survival may be due in part to intrinsic
319 qualities of scarlet macaws. The species lives in a wide
320 range of habitats (Forshaw, 1989) and retains high levels
321 of genetic diversity (Nader et al., 1999). In addition the
322 released birds were only in captivity for 2 generations or
323 less reducing the time for domestication (Wiley et al.,
324 1992). If similar releases were tried with an endangered

325 habitat and diet specialist with low genetic heterozy-
326 gosity the results may not have been as positive (Griffith
327 et al., 1989).

328 Wild predators can rapidly decimate groups of rein-
329 troduced organisms especially when the release candi-
330 dates are captive-raised and lack appropriate anti-
331 predator responses (Snyder et al., 1994; Sinclair et al.,
332 1998). In both Curú and Golfito, the large eagles and
333 hawk-eagles were either extinct or at such low densities
334 as to be irrelevant to the release efforts. Tambopata has
335 5 raptor species large enough to take adult macaws but
336 this did not result in low survivorship possibly due to
337 naïve birds learning from the wild population. Of great
338 relevance to future macaw releases is the fact that avian
339 predators large enough to capture adult macaws occur
340 at naturally low densities and are usually rare or extinct
341 where humans have eliminated macaw populations
342 (Willis and Eisenmann, 1979; Terborgh et al., 1990;
343 Thiollay, 1994; Stotz et al., 1996; BirdLife International,
344 2000). This may allow large macaws to avoid high rates
345 of predation that have plagued reintroductions of
346 smaller Psittacines like Puerto Rican and Thick-billed
347 Parrots (Snyder et al., 1994; USFWS, 2002).

348 Ninety percent of the macaws released for this study
349 were hand-raised. In both Curú and Tambopata the
350 hand-raising was supplemented with frequent human
351 contact. Based on the traditionally poor performance of
352 captive-raised animals, the high survival rates found
353 here were surprising (Griffith et al., 1989; Beck et al.,
354 1994; Snyder et al., 1994). In both Curú and Golfito the
355 birds adapted to life in the wild in the absence of an
356 established wild population (see Lima and Sampaio,
357 2002 for similar results with *Aratinga* parakeets).

As has been found elsewhere, larger releases were more successful than smaller ones (Snyder et al., 1994; Wolf et al., 1998). For this reason future macaw and parrot releases should involve as many birds as is feasible. The number of birds established at the site before release did not significantly correlate with higher survival but anecdotal evidence suggests that this may be important. An eagle killed the first bird released at Tambopata but after the establishment of a core flock, eagle arrivals prompted released birds to alarm call and fly to the lodge for safety (AC pers. obs.). In Golfito the first bird that became established at the release site re-entered the cage where the other macaws were being held. Also at Golfito many birds from the first releases left the site and never returned. After the establishment of the core flock, new releases that left returned accompanying the flocks of established birds. This is similar to the behavior of blue-and-yellow macaws released on Trinidad (Oehler et al., 2001 as corrected by B. Plair pers. com.). Social interactions among Hispanolan Parrots were also important as the presence of birds from earlier releases facilitated the integration of new releases into flocks (Collazo et al., 2003).

Maintaining social interactions among released birds and establishing core flocks appear to be important to the success of parrot releases. Our experience suggests that extended periods of supplemental feeding promoted social interactions among the flock members; encouraged birds to stay in protected areas; allowed project personnel to monitor survival and reproduction; and allowed new releases to quickly find and join the established flock (see also Casimir et al., 2001). Release guidelines for parrots recommend that feeding be conducted only until birds are self-sufficient (Snyder et al., 2000). We suggest that releases continue supplemental feeding even after it is considered superfluous for nutritional reasons. However, care must be taken to ensure that the feeding does not increase predation risk or create birds that approach humans for food (Snyder et al., 1994).

5.2. Reproduction

The key to successful establishment of new populations is reproduction. Successful breeding has taken place only in Tambopata and here hand-raised birds bred with wild mates. In Costa Rica, pairs defended nests (Curú and Golfito) and laid eggs (Golfito). Given the breeding attempts recorded so far there is no a priori reason to think that pairs will not breed at all sites. However, it is unclear if reproduction will be sufficient to allow the populations to grow and expand as hoped. In both Curú and Golfito future releases are scheduled to include more parent raised birds, which may have higher reproductive success (Meyers et al., 1988).

5.3. Raising birds for release

Many hand-raised animals lack the social skills needed to survive and reproduce in the wild (Snyder et al., 1987, 1994; Wiley et al., 1992; Snyder and Snyder, 2000). At all sites our birds formed coherent flocks, formed stable pairs and attempted to breed (see also Sanz and Grajal, 1998; Casimir et al., 2001; Lima and Sampaio, 2002). Our birds probably showed adequate social behavior because they spent significantly more time during the raising process socializing with macaws than with humans (Styles, 2001). In our work the only birds that showed inappropriate social behaviors were confiscated ex-pets who were probably raised in close contact with humans and isolation from conspecifics.

Lack of fear of humans is dangerous as local people often capture or kill released parrots (Snyder et al., 1987; Clubb and Clubb, 1992b; Wiley et al., 1992; Oehler et al., 2001). All birds released in our studies seemed to be more tolerant of humans than wild birds. The Tambopata birds approached humans because they were hand-fed long after weaning age (Table 1). At Curú and Golfito birds did not approach people presumably because weaned birds were kept in cages where they learned to eat food from feeders, not directly from caretakers.

Infectious disease concerns are often cited as the key reason not to conduct releases of captive parrots (Wiley et al., 1992; Snyder et al., 1996). Diagnostic tests for important psittacine diseases such as psittacosis (*Chlamydia psittaci*), avian polyomavirus, and psittacid herpesvirus 1 (Pacheco's disease) may not always detect these agents. Additionally, diseases such as proventricular dilatation disease (wasting disease) are of unknown etiology and no diagnostic test exists. Unfortunately these diseases may remain sub-clinical until stress precipitates an active infection or a susceptible host is infected after contact (Altman et al., 1997; D. Styles pers. com.). As a result, appropriate biosecurity measures should be instituted to ensure that release candidates are protected from infectious disease. In Tambopata, the most remote of the three release sites, hand-raised birds contracted *Salmonella* during the rearing process (Kareesh et al., 1997). Additionally at Curú, the one bird that tested positive for psittacosis was not raised at the Amigos de las Aves facility. This emphasizes the potential risk of using birds from insecure sources and reaffirms that all release programs must have strict quarantine, biosecurity, and disease testing regimens to ensure the production of disease free release candidates (Snyder et al., 1996). The threat of spreading infectious diseases from captive to wild populations exists and therefore releases should not be conducted in areas with viable populations of wild conspecifics.

The release of ex-pet parrots in to the wild is often considered by conservation-minded pet-owners. Our

467 experience shows that ex-pets are the worst candidates
 468 for release due to their failure to interact appropriately
 469 with other macaws and their propensity to stay near
 470 humans. In addition confiscated birds, be they wild
 471 caught or ex-pets, are always a disease risk. Birds that
 472 are poached or held in homes are often kept in poor
 473 conditions, fed improper diets and exposed to other
 474 captive wild birds or domestic fowl (Nilsson, 1981).
 475 These are ideal conditions for the development of seri-
 476 ous diseases.

477 The results from these case studies show that properly
 478 socialized hand-raised scarlet macaws survive in the
 479 wild. The high survival rates found here may be due to
 480 the innate adaptability of scarlet macaws and inherently
 481 low predation rates on these large birds. No pairs of
 482 released captive bred birds have reproduced successfully
 483 so it is uncertain if these populations will become self-
 484 sustaining. While these results may not be duplicable
 485 with all species, the current study shows that captive
 486 breeding and reintroduction can be used to reestablish
 487 psittacines in areas from which they have been extir-
 488 pated.

489 Acknowledgements

490 We thank J. I. Rojas, L. Zapater, E. Nycander and all
 491 the project assistants. Thanks to the Ministerio de
 492 Ambiente y Energía (Costa Rica) and the Instituto
 493 Nacional de Recursos Naturales (Peru). Funding pro-
 494 vided by The EarthWatch Institute, Wildlife Conserva-
 495 tion International, Rainforest Expeditions, Kaytee
 496 Avian Foundation, Chiquita, Raleigh-Durham Caged
 497 Bird Society and private donors and editorial comments
 498 by N. Snyder, K. Renton, D. Homberger, D. Styles, G.
 499 Matuzak, J. Massello, and E. Villalobos.

500 References

501 Altman, R.B., Clubb, S.L., Dorrestien, G.M., Quesenberry, K., 1997.
 502 Avian Medicine and Surgery. W.B. Saunders, Philadelphia, Penn-
 503 sylvania.
 504 Balmford, A., Mace, G.M., Leader-Williams, N., 1996. Designing the
 505 ark: setting priorities for captive breeding. *Conservation Biology*
 506 10, 719–727.
 507 Beck, B.B., Rapaport, L.G., Price, M.R.S., Wilson, A.C., 1994.
 508 Reintroduction of captive-born animals. In: Olney, P.J.S., Mace,
 509 G.M., Feistner, A.T.C. (Eds.), *Creative Conservation: Interactive*
 510 *Management of Wild and Captive Animals*. Chapman & Hall,
 511 London, pp. 265–286.
 512 Bennett, P.M., Owens, I.P.F., 1997. Variation in extinction risk among
 513 birds: chance or evolutionary predisposition? *Proceedings of the*
 514 *Royal Society of London B* 264, 401–408.
 515 Biggins, D.E., Vargas, A., Godbey, J.L., Anderson, S.H., 1999.
 516 Influence of prerelease experience on reintroduced black-footed
 517 ferrets (*Mustela nigripes*). *Biological Conservation* 89, 121–129.
 518 BirdLife International, B., 2000. *Threatened birds of the world*. Lynx
 519 Edicions and BirdLife International, Barcelona and Cambridge.

Bjork, R., Powell, G., 1995. Buffon's Macaw: some observations on 520
 the Costa Rican populations, its lowland forest habitat and 521
 conservation. In: Abramson, J., Spear, B.L., Thomsen, J.B. 522
 (Eds.), *The Large Macaws: Their Care, Breeding and Conserva-* 523
tion. Raintree Publications, Ft. Bragg, CA, pp. 387–392. 524
 Brightsmith, D.J. Effects of weather on avian geophagy in Tambopata, 525
 Peru. *Wilson Bulletin* (in press). 526
 Casimir, D., Fisher, S., Rodda, D., Nichols, R., Morris, A., Woolaver, 527
 L., 2001. The release of captive-raised echo parakeets to the wild, 528
 2000/2001. Unpublished report, Mauritian Wildlife Foundation, 529
 Mauritius. 530
 Clubb, S.L., 1992. The role of private aviculture in the conservation of 531
 neotropical psittacines. In: Snyder, N.F.R., Beissinger, S.R. (Eds.), 532
New World Parrots in Crisis. Smithsonian Institution Press, 533
 Washington, DC, pp. 117–132. 534
 Clubb, K.J., Clubb, S.L., 1992a. Status of macaws in aviculture 535
 (Chapter 24). In: Shubot, R.M., Clubb, K.J., Clubb, S.L. (Eds.), 536
Psittacine Aviculture: Perspectives, Techniques and 537
Research. Avicultural Breeding and Research Center, Loxahatchee, 538
 FL. 539
 Clubb, K.J., Clubb, S.L., 1992b. Reintroduction of military macaws in 540
 Guatemala (Chapter 23). In: Shubot, R.M., Clubb, K.J., Clubb, 541
 S.L. (Eds.), *Psittacine Aviculture: Perspectives, Techniques and* 542
Research. Avicultural Breeding and Research Center, Loxahatchee, 543
 FL. 544
 Collar, N.J., 1997. Family psittacidae. In: Hoyo, J.d., Elliott, A., 545
 Sargatal, J. (Eds.), *Handbook of the Birds of the World*. Lynx 546
 Edicions, Barcelona, Spain, pp. 280–479. 547
 Collazo, J.A., White, T.H., Vilella, F.J., Guerrero, S., 2003. Survival of 548
 captive-reared Hispaniolan Parrots released in Parque Nacional del 549
 Este, Dominican Republic. *Condor* 105, 198–207. 550
 Cox, D.R., Snell, E.J., 1989. *Analysis of Binary Data*. Chapman & 551
 Hall, New York. 552
 Derrickson, S.R., Snyder, N.F.R., 1992. Potentials and limits of 553
 captive breeding in parrot conservation. In: Snyder, N.F.R., 554
 Beissinger, S.R. (Eds.), *New World Parrots in Crisis*. Smithsonian 555
 Institution Press, Washington, DC, pp. 133–164. 556
 Forshaw, J.M., 1989. *Parrots of the World*, third ed. Landsdowne 557
 Editions, Melbourne, Australia. 558
 Foster, R.B., Parker, T.A., Gentry, A.H., Emmons, L.H., Chicchón, 559
 A., Schulenberg, T., Rodríguez, L., Larnas, G., Ortega, H., 560
 Icochea, J., Wust, W., Romo, M., Alban, C.J., Phillips, O., Reynel, 561
 C., Kratter, A., Donahue, P.K., Barkley, L.J., 1994. The Tambo- 562
 pata–Candamo reserved zone of southeastern Peru: a biological 563
 assessment. *Conservation International*, Washington, DC. Avail- 564
 able from: <http://www.biodiversityscience.org/xp/CABS/publications/cabs_publ_research/rap_bulletins/terrestrial_rap_bulletins/terrappbulletins.xml> last accessed, April 2004. 565
 Gippoliti, S., Carpaneto, G.M., 1997. Captive breeding, zoos and good 566
 sense. *Conservation Biology* 11, 806–807. 567
 Griffith, B., Scott, J.M., Carpenter, J.W., Reed, C., 1989. Transloca- 568
 tion as a species conservation tool: status and strategy. *Science* 245, 569
 477–480. 570
 Griscom, B.W., Ashton, P.M.S., 2003. Bamboo control of forest 571
 succession: *Guadua sarcocarpa* in Southeastern Peru. *Forest Ecol-* 572
ogy and Management 175, 445–454. 573
 Holdridge, L.R., 1967. *Life Zone Ecology*, Revised Edition. Tropical 574
 Science Center, San Jose, Costa Rica. 575
 Hensler, G.L., Nichols, J.D., 1981. Mayfield method of estimating 576
 nesting success: a model, estimators and simulation results. *Wilson* 577
Bulletin 93, 42–53. 578
 Janik, D., Ramirez, S., Fournier, R., Sibaja, R., Guillén, F., 2003. 579
 Informe del Proyecto de Reintroducción de la Lapa Roja (*Ara* 580
macao) en Playa San Josecito, Golfito. Fundación Restauración de 581
 la Naturaleza, San Jose, Costa Rica. 582
 Johnson, K.A., 1992. 1991 Psittacine Captive Breeding Survey. 583
 TRAFFIC (USA), Washington, DC. 584
 585
 586

- 587 Juniper, T., 2003. Spix's Macaw: The Race to Save the World's Rarest
588 Bird. Fourth Estate, London.
- 589 Juniper, T., Parr, M., 1998. Parrots: A Guide to Parrots of the World.
590 Yale University Press, New Haven.
- 591 Karesh, W.B., Campo, A.d., Braselton, W.E., Puche, H., Cook, R.A.,
592 1997. Health evaluation of free-ranging and hand-reared macaws
593 (*Ara* spp.) in Peru. *Journal of Zoo and Wildlife Medicine* 28, 368–
594 377.
- 595 Lima, P.C., Sampaio, S., 2002. Reintroduction of captive-bred gold-
596 capped conures in Bahia. *PsittScene* 53, 4–5.
- 597 Mayfield, H.F., 1975. Suggestions for calculating nest success. *Wilson*
598 *Bulletin* 87, 456–466.
- 599 Meyers, S.A., Millam, J.R., Roudybush, T.E., Grau, C.R., 1988.
600 Reproductive success of hand-reared vs. parent-reared Cockatiels.
601 *Auk* 105, 536–542.
- 602 Nader, W., Werner, D., Wink, M., 1999. Genetic diversity of scarlet
603 macaws *Ara macao* in reintroduction studies for threatened
604 populations in Costa Rica. *Biological Conservation* 87, 269–272.
- 605 Nilsson, G., 1981. The bird business: A Study of the Commercial Cage
606 Bird Trade, second ed. Animal Welfare Institute, Washington, DC.
- 607 Noss, R., 2001. Why restore large mammals. In: Maehr, D., Noss, R.,
608 Larkin, J. (Eds.), *Large Mammal Restoration. Ecological and*
609 *sociological challenges in the 21st century*. Island Press, pp. 1–21.
- 610 Nycander, E., Blanco, D.H., Holle, K.M., Campo, A.d., Munn, C.A.,
611 Moscoso, J.I., Ricalde, D.G., 1995. Manu and Tambopata: nesting
612 success and techniques for increasing reproduction in wild macaws
613 in southeastern Peru. In: Abramson, J., Spear, B.L., Thomsen, J.B.
614 (Eds.), *The Large Macaws: Their Care, Breeding and Conserva-*
615 *tion*. Raintree Publications, Ft. Bragg, CA, pp. 423–443.
- 616 Oehler, D.A., Boodoo, D., Blair, B., Kuchinski, K., Campbell, M.,
617 Lutchmendial, G., Ramsbidge, S., Maruska, E.J., Malowski, S.,
618 2001. Translocation of blue-and-gold macaw *Ara ararauna* into its
619 historical range on Trinidad. *Bird Conservation International* 11,
620 129–141.
- 621 Pyle, P., Nur, N., Henderson, R.P., Desante, D.F., 1993. The effects of
622 weather and lunar cycle on nocturnal migration of landbirds at
623 southeast Farallon Island, California. *Condor* 95, 343–361.
- 624 Renton, K., 2000. Scarlet macaw. In: Reading, R.P., Miller, B. (Eds.),
625 *Endangered Animals: A Reference Guide to Conflicting Issues*.
626 Greenwood Press, Westport, Connecticut.
- 627 Sanz, V., Grajal, A., 1998. Successful reintroduction of captive-raised
628 yellow-shouldered amazon parrots on Margarita Island, Venezuela.
629 *Conservation Biology* 12, 430–441.
- 630 Schutt, A., Vaughan, C., 1995. Incorporating wildlife into develop-
631 ment: The case of the Curú wildlife refuge and farm, Costa Rica.
632 In: Bissonette, J.A., Krausman, P.R., (Eds.), *Integrating People*
633 *and Wildlife for a Sustainable Future. Proceedings of the First*
634 *International Wildlife Management Congress*, pp. 250–254.
- 635 Sinclair, A.R.E., Pech, R.P., Dickman, C.R., Hik, D., Mahon, P.,
636 Newsome, A.E., 1998. Predicting effects of predation on conserva-
637 tion of endangered prey. *Conservation Biology* 12, 564–575.
- Snyder, N.F.R., Derrickson, S.R., Beissinger, S.R., Wiley, J.W., 638
Smith, T.B., Toone, W.D., Miller, B., 1996. Limitations of captive 639
breeding in endangered species recovery. *Conservation Biology* 10, 640
338–348. 641
- Snyder, N.F.R., Derrickson, S.R., Beissinger, S.R., Wiley, J.W., 642
Smith, T.B., Toone, W.D., Miller, B., 1997. Limitations of captive 643
breeding: reply to Gippoliti and Carpaneto. *Conservation Biology* 644
11, 808–810. 645
- Snyder, N.F.R., Koenig, S.E., Koschmann, J., Koschmann, J., Snyder, 646
H.A., Johnson, T.B., 1994. Thick-billed parrot releases in Arizona. 647
Condor 96, 845–862. 648
- Snyder, N.F.R., McGowan, P., Gilardi, J., Grajal, A., 2000. Parrots. 649
Status survey and conservation action plan 2000–2004. IUCN, 650
Gland, Switzerland and Cambridge, UK, pp. 180. 651
- Snyder, N.F.R., Snyder, H., 2000. *The California Condor: A Saga of* 652
Natural History and Conservation. Academic Press, New York. 653
- Snyder, N.F.R., Wiley, J.W., Kepler, C.B., 1987. *The Parrots of* 654
Luquillo: Natural History and Conservation of the Puerto Rican 655
parrot. Western Foundation of Vertebrate Zoology, Los Angeles, 656
CA. 657
- Sokal, R.R., Rohlf, F.J., 1995. *Biometry*. Freeman, New York. 658
- Stotz, D.F., Fitzpatrick, J.W., Parker III, T.A., Moskovits, D.K., 659
1996. *Neotropical Birds: Ecology and Conservation*. University of 660
Chicago Press, Chicago. 661
- Styles, D.K., 2001. Captive psittacine behavioral reproductive hus- 662
bandry and management: socialization, aggression control, and 663
pairing techniques. In: *The Association of Avian Veterinarians* 664
2001 Conference Proceedings. The Association of Avian Veteri- 665
narians, Orlando, FL, pp. 3–14. 666
- Terborgh, J., Robinson, S.K., Parker III, T.A., Munn, C., Pierpont, 667
N., 1990. Structure and organization of an Amazonian forest bird 668
community. *Ecological Monographs* 60, 213–238. 669
- Thiollay, J.-M., 1994. Structure, density and rarity in an Amazonian 670
rainforest bird community. *Journal of Tropical Ecology* 10, 449– 671
481. 672
- USFWS, 2002. Puerto Rican Parrot 2002 update, Washington, DC. 673
Available from: <[http://southeast.fws.gov/pubs/facts/pr_par-](http://southeast.fws.gov/pubs/facts/pr_parrot_update.pdf) 674
[rot_update.pdf](http://southeast.fws.gov/pubs/facts/pr_parrot_update.pdf)>. 675
- Wiedenfeld, D.A., 1994. A new subspecies of scarlet macaw and its 676
status and conservation. *Ornitologia Neotropical* 5, 99–104. 677
- Wiley, J.W., Snyder, N.F.R., Gnam, R.S., 1992. Reintroduction as a 678
conservation strategy for parrots. In: Beissinger, S.R., Snyder, 679
N.F.R. (Eds.), *New World Parrots in Crisis*. Smithsonian Institu- 680
tion Press, Washington, DC, pp. 165–200. 681
- Willis, E.O., Eisenmann, E., 1979. A revised list of the birds of Barro 682
Colorado Island, Panama. Smithsonian Institution Press, Wash- 683
ington, DC. 684
- Wolf, C.M., Garland, T., Griffith, B., 1998. Predictors of avian and 685
mammalian translocation success: reanalysis with phylogeneti- 686
cally independent contrasts. *Biological Conservation* 86, 243– 687
255. 688